

Hamiltonian Paths Through Two- and Three-Dimensional Grids

Volume 110

Number 2

March-April 2005

William F. Mitchell

National Institute of Standards
and Technology,
Gaithersburg, MD 20899-8910

william.mitchell@nist.gov

This paper addresses the existence of Hamiltonian paths and cycles in two-dimensional grids consisting of triangles or quadrilaterals, and three-dimensional grids consisting of tetrahedra or hexahedra. The paths and cycles may be constrained to pass from one element to the next through an edge, through a vertex, or be unconstrained and pass through either. It was previously known that an unconstrained Hamiltonian path exists in a triangular grid under very mild conditions, and that there are triangular grids for which there is no through-edge Hamiltonian path. In this paper we prove that a through-vertex Hamiltonian cycle exists in any triangular or tetrahedral grid

under very mild conditions, and that there exist quadrilateral and hexahedral grids for which no unconstrained Hamiltonian path exists. The existence proofs are constructive, and lead to an efficient algorithm for finding a through-vertex Hamiltonian cycle.

Key words: adaptive grid refinement; Hamiltonian path; load balancing; refinement-tree partition.

Accepted: February 23, 2005

Available online: <http://www.nist.gov/jres>

1. Introduction

Adaptive grid refinement has been shown to be an effective tool for reducing the size of the grid (and consequently the linear system) required for a given accuracy when numerically solving partial differential equations. Problems involving singularities or multi-scale behavior practically require adaptive refinement. When implemented for parallel computers, dynamic load balancing is required to keep all of the processors busy. This involves partitioning the grid into equal sized pieces (in some measure), and distributing the data among the processors accordingly. Many of the methods for computing a partition are based on a linearization of the elements (or path through the elements) in a two-dimensional or three-dimensional grid, and then cutting the linear sequence into pieces of equal size. Some of the methods that fall into this category are the space filling curves [1], OCTREE [2], and refinement-tree [3] methods. Further information on partitioning methods can be found in Ref. [4].

The space filling curve and OCTREE methods are not guaranteed to create connected partitions, which may be a desirable property. The refinement-tree method is guaranteed to give connected partitions provided that an appropriate linearization of the initial coarse grid is given. Such a linearization would be a Hamiltonian path, i.e., a path which passes from an element to a neighboring element and goes through each element exactly once. Heber et al. [5] proved that, for grids consisting of triangles, a Hamiltonian path always exists under very mild conditions. Moreover, the proof is a constructive proof which leads to an efficient algorithm to find a Hamiltonian path. However, the refinement-tree partitioning algorithm requires a stronger result in which the Hamiltonian path always passes through vertices when moving from one element to the next (as opposed to passing through edges), and does not go out the same vertex through which it came in. We call this a through-vertex Hamiltonian path. Heber's algorithm produces Hamiltonian paths that pass through both vertices and

edges. Additionally, Ref. [5] only addresses triangles, not the other elements considered in this paper.

The main result of this paper is the proof that there always exists a through-vertex Hamiltonian path in grids consisting of triangles or tetrahedra, under very mild conditions. The proof is constructive, which leads to an algorithm to construct such a path. We do not explicitly give the algorithm in this paper, but it follows easily from the proofs of the main theorems. Little is known about the conditions under which a Hamiltonian path exists in grids consisting of quadrilaterals or hexahedra. An algorithm is given that might find a through-vertex Hamiltonian path in a quadrilateral or hexahedral grid, if one exists, and is likely to give a broken path with a small number of discontinuities, i.e., something close to a through-vertex Hamiltonian path.

The remainder of the paper is organized as follows. In Sec. 2 we introduce the notation and define the terms used in this paper. Section 3 addresses triangles and tetrahedra. It reviews previously known results and presents the main results of this paper. In Sec. 4 we discuss quadrilaterals and hexahedra. Section 5 contains the conclusions.

2. Definitions

For the purposes of this paper, an *element*, E , is a triangle, quadrilateral, tetrahedron, or hexahedron. An element contains *vertices*, v , and *edges*, e , with the obvious definitions, and three-dimensional elements contain *faces*, f .

A *grid*, G , is the union of a collection of elements, $\{E_i\}$, all of the same kind, such that $G = \cup E_i$ is a connected, bounded region in \mathbb{R}^2 or \mathbb{R}^3 , and $E_i \cap E_j = \phi$, $i \neq j$, where $\overset{\circ}{E}$ denotes the interior of element E , and ϕ is the empty set. We say that E_i is an element of G , $E_i \in G$. A vertex of E_i is a *boundary vertex* if it lies on the boundary of G , and an *interior vertex* if it is not a boundary vertex. We say the size of G , $|G|$, is N if there are N elements in G . A *triangular grid*, *quadrilateral grid*, *tetrahedral grid*, and *hexahedral grid* is a grid consisting entirely of triangles, quadrilaterals, tetrahedra and hexahedra, respectively.

A grid is said to be conforming if $E_i \cap E_j$, $i \neq j$, is a common vertex, common edge, common face or empty. A vertex of an element is called a *hanging node* if it lies in the interior of an edge or face of another element. It follows immediately from the definition that a conforming grid has no hanging nodes. A grid is said to be

1-nonconforming if there is at least one hanging node in the grid, all edges contain at most one hanging node, the interior of all faces contain at most one hanging node, and the intersection of two elements is a vertex, edge or face of one of the elements, or empty. See Fig. 1 for examples of conforming and 1-nonconforming grids. This paper will primarily consider triangular and tetrahedral grids that are conforming, and quadrilateral and hexahedral grids that are conforming or 1-nonconforming.

A *path* with *length* n in a grid G is a sequence of elements, $E_1 E_2 \dots E_n$, $E_i \in G$, $i = 1, n$, with $E_i \cap E_{i+1} \neq \phi$, and $E_i \neq E_{i+1}$, $i = 1, n-1$. A *cycle* of length n is a path of length $n+1$ in which $E_1 = E_{n+1}$. A *Hamiltonian path* is a path in which every element in G appears exactly once. A *Hamiltonian cycle* is a cycle in which every element in G appears exactly once except for $E_1 = E_{n+1}$, which appears exactly twice.

A sequence of elements $E_1 E_2 \dots E_n$ for which there exists an i with $E_i \cap E_{i+1} = \phi$ is called a *broken path*, and the sequence $E_i E_{i+1}$ is called a *discontinuity* in the path.

A *through-vertex path* (*through-vertex cycle*) is a path (cycle) in which the passage from one element to the next is specified as a common vertex, and the path does not pass through the same vertex when entering and exiting an element. Specifically, it is $E_1 v_1 E_2 v_2 \dots v_{n-1} E_n$ where $v_i \subseteq E_i \cap E_{i+1}$, $i = 1, n-1$, $v_i \neq v_{i+1}$, $i = 1, n-2$, and $E_1 E_2 \dots E_n$ is a path (cycle). We say that the path *passes through* v_i when going from E_i to E_{i+1} , and that we come into E_i through the *in-vertex* v_{i-1} and leave E_i through the *out-vertex* v_i . A *through-edge* path and cycle, and *through-face* path and cycle are defined similarly. A *through-vertex Hamiltonian path* is a through-vertex path that is also a Hamiltonian path, and the definitions of the obvious similar terms are similar. An *unconstrained path* is one that may pass through any of vertices, edges or faces. Although the term is redundant, we will use it when we want to emphasize that the path is not constrained to pass through only vertices, edges or faces.

A vertex, v , is called a *cut vertex* if $G \setminus v$ is disconnected. See Fig. 2. A *local cut vertex* is a vertex whose removal causes G to become disconnected locally. Formally, v is a local cut vertex if there exists an $R > 0$ such that for all $0 < \epsilon < R$, $(G \cap B(v, \epsilon)) \setminus v$ is disconnected, where $B(v, \epsilon)$ is the ball of radius ϵ centered at v . Note that a cut vertex is also a local cut vertex. Figure 2(b) illustrates a local cut vertex that is not a cut vertex. In three dimensions, the terms *cut edge* and *local cut edge* are defined similarly.




Fig. 1. (a) A conforming triangular grid. (b) A 1-nonconforming quadrilateral grid. (c) A grid that is neither conforming nor 1-nonconforming, because an element edge contains more than one hanging node.




Fig. 2. (a) A grid containing a cut vertex. (b) A grid containing a local cut vertex. In each case, the vertex is shown as a large dot.

3. Triangles and Tetrahedra

In this section we present what is known about the existence of the different types of Hamiltonian paths and cycles for conforming triangular and tetrahedral grids. We begin with a review of previously published results. We then give the main results of this paper, which are the existence of through-vertex Hamiltonian cycles under very mild conditions. Counterexamples are presented throughout to show conditions under which Hamiltonian cycles and paths do not exist, and to show that the conditions in the hypotheses of the theorems are essential.

Note that the existence of a cycle implies the existence of a path. Therefore most of the existence statements are made for cycles, and it is understood that the same statement holds for paths.

Conversely, non-existence statements are usually made about paths, and it is understood that the same statement holds for cycles. The exceptions to this are when quoting statements from other papers and when a path exists but a cycle does not.

The following theorem is due to Heber et al. [5].

Theorem 1 *There exists a Hamiltonian path for any conforming triangular grid that contains no local cut vertices.*

The statement of the theorem in Ref. [5] does not mention the absence of local cut vertices or that the grid must be conforming, however the definition of a grid in that paper is that it be a “simplicial complex coming from the simplicial decomposition of a connected 2D

manifold” which implies these conditions. The theorem also holds for Hamiltonian cycles, simply by starting the base case in Heber’s inductive proof with a Hamiltonian cycle between two triangles. The hypotheses of the theorem need to be extended slightly because the definition of a cycle assumes at least two elements.

Corollary 1 *There exists a Hamiltonian cycle for any conforming triangular grid, of size at least 2, that contains no local cut vertices.*

Hamiltonian paths and cycles also exist for tetrahedral grids under similar conditions. This follows immediately from Theorem 2 later in this section.

The following counterexamples show that the absence of cut vertices is an essential condition for Theorem 1, and the absence of local cut vertices is an essential condition for Corollary 1. However, it is not known whether the absence of local cut vertices is essential for the Hamiltonian path.

Counterexample 1 *There exists a triangular grid containing cut vertices for which there is no Hamiltonian path. See Fig. 3.*

Counterexample 2 *There exists a triangular grid containing local cut vertices (but no cut vertices) for which there is no Hamiltonian cycle. See Fig. 4.*

Corollary 1 says that, under very mild conditions, we can always find a Hamiltonian cycle (and hence a Hamiltonian path) in a triangular grid. This is an unconstrained Hamiltonian cycle, i.e., it does not say whether the passage from one element to the next is through a vertex or edge. Indeed, the recursive algorithm implied by the proof of Theorem 1 in Ref. [5] will usually result in passages through both vertices and edges. The obvious question is whether or not through-vertex Hamiltonian cycles or paths and through-edge Hamiltonian cycles or paths exist. The following well-known counterexample shows that we cannot expect to find a through-edge Hamiltonian path in a triangular grid. In fact, determining whether or not a through-edge Hamiltonian cycle exists in a triangular grid is known to be NP-complete [6]. A similar counterexample can be constructed for through-face Hamiltonian paths in tetrahedral grids.

Counterexample 3 *There exists a conforming triangular grid with no local cut vertices for which there is no through-edge Hamiltonian path. See Fig. 5.*




Fig. 3. Example of a triangular grid containing cut vertices for which there is no Hamiltonian path.




Fig. 4. Example of a triangular grid containing local cut vertices for which there is no Hamiltonian cycle.




Fig. 5. Example of a conforming triangular grid without local cut vertices for which there is no through-edge Hamiltonian path.

The situation is less grim for through-vertex Hamiltonian cycles. The main result of this paper is that through-vertex Hamiltonian cycles exist for triangular and tetrahedral grids under conditions similar to those for the existence of an unconstrained Hamiltonian cycle. They again require a conforming grid with no local cut vertices. Triangular grids also require that there be at least one interior vertex, and tetrahedral grids also require that there be no local cut edges.

The following lemma says that, under conditions that will arise in the proofs of the main theorems, we can always find two triangles that share a edge, or two tetrahedra that share a face.

Lemma 1 *Let G be a conforming triangular (tetrahedral) grid with $|G| \geq 2$, no local cut vertices and no local cut edges. Let $G_1 \subset G$ contain no local cut vertices and no local cut edges, $G_2 = G \setminus G_1$, $|G_1| \geq 1$, and $|G_2| \geq 1$. Then*

1. *there exists $E_1, E_2 \in G$ such that $E_1 \cap E_2$ is a common edge (face), and*
2. *there exists $E_1 \in G_1$ and $E_2 \in G_2$ such that $E_1 \cap E_2$ is a common edge (face).*

Proof: For part 1, suppose there are no elements that share an edge (face). Then all connections between elements are vertices (or edges), which must then be local cut vertices (or local cut edges) contradicting the hypothesis that there are no local cut vertices or local cut edges.

For part 2, suppose there is no element in G_2 that shares an edge (face) with an element in G_1 . Then G_1 and G_2 are connected by only vertices (and edges), which must then be local cut vertices (or local cut edges). \square

We first give the main result for tetrahedral grids, where the proof is shorter.

Theorem 2 *Let G be a conforming tetrahedral grid with $|G| \geq 2$. If G contains no local cut vertices and no local cut edges then there exists a through-vertex Hamiltonian cycle for G .*

Proof: We prove this by induction on $|G_1|$ where $G_1 \subseteq G$, G_1 satisfies the hypotheses of the theorem, and we can exhibit a through-vertex Hamiltonian cycle for G_1 .

For $|G_1| = 2$, let $G_1 = \{E_1, E_2\}$ where E_1 and E_2 are any two elements that share a common face. Lemma 1 insures the existence of these elements. Let v_1 and v_2 be two of the vertices that they share. Then $E_1 v_1 E_2 v_2 E_1$ is a through-vertex Hamiltonian cycle for this subgrid.

By induction, let $k = |G_1|$ and suppose we have $G_1 \subset G$, $k \geq 2$, G_1 satisfies the hypotheses of the theorem, and $E_1 v_1 E_2 v_2 \dots E_k v_k E_1$ is a through-vertex Hamiltonian cycle for G_1 . The grid and cycle can be extended to size $k + 1$ as follows.

Let $E_{k+1} \in G \setminus G_1$ be an element that shares a face with some element $E_i \in G_1$. The existence of E_{k+1} is guaranteed by Lemma 1. Without loss of generality, assume E_i is not E_1 (otherwise, just start the numbering of the cycle with a different element). Since E_i has only four vertices, one of the three vertices shared by E_i and

E_{k+1} must be either E_i 's in-vertex v_{i-1} or E_i 's out-vertex v_i . Without loss of generality, suppose it is v_i (otherwise, just reverse the ordering of the cycle). One of the three shared vertices, say v , must not be v_{i-1} or v_i . Then a through-vertex Hamiltonian cycle in $G_1 \cup E_{k+1}$ is $E_1 \dots v_{i-1} E_i v E_{k+1} v_i E_{i+1} \dots E_1$. \square

The proof for triangular grids is also a constructive proof that begins with a cycle through two elements and inductively extends this cycle to the complete grid. However, it is a more complicated proof because, in the notation of Theorem 2, we cannot guarantee that E_{k+1} shares a vertex with E_i that is not v_{i-1} or v_i . In that case, a more complicated extension of the cycle must be performed.

Theorem 3 *Let G be a conforming triangular grid with $|G| \geq 2$. If G contains no local cut vertices and has at least one interior vertex then there exists a through-vertex Hamiltonian cycle for G .*

Proof: As in Theorem 2, we prove this by induction on $|G_1|$, and the base case with two triangles is trivial. By induction, suppose we have $G_1 \subset G$, $|G_1| = k \geq 2$, G_1 contains no local cut vertices, and $E_1 v_1 E_2 v_2 \dots E_k v_k E_1$ is a through-vertex Hamiltonian cycle for G_1 . The grid and cycle can be extended to a larger size as follows.

Let $E_{k+1} \in G \setminus G_1$ be an element that shares an edge with some element $E_i \in G_1$. The existence of E_{k+1} is guaranteed by Lemma 1. Without loss of generality, assume E_i is not E_1 (otherwise, just start the numbering of the cycle with a different element). One of the two vertices that E_{k+1} shares with E_i must be either E_i 's in-vertex, v_{i-1} , or E_i 's out-vertex, v_i . Without loss of generality, assume it shares the out-vertex (otherwise, just reverse the order of the cycle). There are four cases to consider.

Case 1. E_{k+1} does not contain v_{i-1} . This is the easy case that corresponds to the proof of Theorem 2. Let v be the other vertex shared by E_{k+1} and E_i . Then the new cycle is $E_1 \dots v_{i-1} E_i v E_{k+1} v_i E_{i+1} \dots E_1$. This extension is illustrated in Fig. 6. The arrows pointing from a vertex to the interior of a triangle or from the interior of a triangle to a vertex denote the in-vertex and out-vertex, respectively.

Case 2. E_{k+1} contains v_{i-1} and there is another triangle, E_{k+2} , not on the current cycle, that shares an edge and v_i with E_{k+1} . This case is illustrated in Fig. 7. Let v be the other vertex shared by E_{k+1} and E_{k+2} . Then the new cycle is $E_1 \dots v_{i-1} E_i v_i E_{k+1} v E_{k+2} v_i E_{i+1} \dots E_1$.

Case 3. E_{k+1} contains v_{i-1} and there is another

triangle, E_{k+2} , that is on the current cycle and shares an edge and v_i with E_{k+1} .

First note that E_{k+1} must contain both the in-vertex and out-vertex of E_{k+2} , otherwise we could apply case 1 with E_{k+2} as E_i to add E_{k+1} to the cycle. Also note that whenever a triangle not on the cycle contains both the in-vertex and out-vertex of a triangle that is on the cycle, we can "swap" this triangle with the other triangle by removing the other triangle from the cycle and inserting the new triangle in its place, as illustrated in Fig. 8.




Fig. 6. Extension of the cycle for case 1 in Theorem 3.




Fig. 7. Extension of the cycle for case 2 in Theorem 3.




Fig. 8. Swapping element E_{k+1} for element E_{k+2} in the cycle.

Case 3a. v_i is an interior vertex.
Swap elements around v_i until either

1. there are two adjacent elements that are not in the cycle (then apply case 2), or
2. there are adjacent elements, one on the cycle and one not on the cycle, where the one not on the cycle does not contain both the in-vertex and out-vertex of the one on the cycle (then apply case 1).

Note that if we do not encounter two adjacent elements that are not in the cycle (the first subcase), then we must eventually reach the second subcase because the other triangle adjacent to E_i cannot contain the in-vertex of E_i , v_{i-1} . See Fig. 9.

Case 3b. v_i is a boundary vertex.
This implies that all three vertices of E_{k+1} are on the boundary, for if not we could select a different E_i , v_{i-1} and v_i (possibly reversing the order of the cycle) such that an interior vertex of E_{k+1} is v_i . This leads to a natural decomposition of G into three components plus E_{k+1} as illustrated in Fig. 10. The intersection of any two components is a vertex of E_{k+1} . Each component contains a triangle that shares an edge with E_{k+1} because there are no local cut vertices. One component contains E_i and a second component contains E_{k+2} . If the third component is not empty, let E be the element that shares an edge with E_{k+1} . If E is not on the cycle, then apply case 2 (reversing the order of the cycle if necessary). If E is on the cycle and E_{k+1} does not contain both the in-vertex and out-vertex of E , then apply case 1 with E as E_i . Thus we can assume that E_{k+1} contains the in-vertex and out-vertex of all adjacent triangles, so it can be swapped with any of them.

Pick an interior vertex of G , v_{int} . Swap E_{k+1} with the neighbor that is in the same component as v_{int} . Apply case 1, 2 or 3a to reinsert the other element, if appropriate. If not, then consider the decomposition around the

new element and repeat the process (see Fig. 11). The component that contains v_{int} has at least lost the swapped triangle, and thus is smaller in this decomposition. Since the size of the component containing v_{int} will continue to shrink with each application of this process, it must eventually lead to v_{int} , at which point case 3a applies, unless it ends earlier by applying case 1, 2 or 3a.

Case 4. E_{k+1} contains v_{i-1} and there is no other triangle that shares an edge and v_i with E_{k+1} , i.e., that edge is on the boundary.

Then there must be another triangle that shares v_i and an edge with E_i , for if that edge of E_i was also on the boundary, then v_i would be a local cut vertex. Swap E_{k+1} with E_i , and then apply either case 2 or case 3 to add E_i back into the path. \square

The inclusion of an interior vertex is an essential condition for Theorem 3. The same counterexample can be used as was used in Counterexample 3. However, this example does contain a through-vertex Hamiltonian path. It is not known whether or not the inclusion of an interior vertex is essential for the existence of a through-vertex Hamiltonian path.

Counterexample 4 *There exists a conforming triangular grid with no local cut vertices and no interior vertices for which there is no through-vertex Hamiltonian cycle. See Fig. 5.*

Since the inclusion of an interior vertex “fixes” Counterexample 4, a natural question is whether it can also “fix” Counterexample 3. The following counterexample says that this is not the case.

Counterexample 5 *There exists a conforming triangular grid with no local cut vertices and at least one interior vertex for which there is no through-edge Hamiltonian path. See Fig. 12.*




Fig. 9. A case where swapping around an interior vertex v_i continues until the other triangle adjacent to E_i is reached.




Fig. 10. Decomposition of G into three components plus E_{k+1} when all three vertices of E_{k+1} are on the boundary.




Fig. 11. Second decomposition of G into three components plus a triangle after swapping E_{k+1} with a neighbor.

4. Quadrilaterals and Hexahedra

It is not known under what conditions any kind of Hamiltonian path or cycle is guaranteed to exist for quadrilateral and hexahedral grids, or even if there is any characterization. It would certainly be much more stringent conditions than for triangles and tetrahedra, as the following counterexample shows. By replacing the squares with cubes, the same counterexample works for hexahedral grids.




Fig. 12. Example of a conforming triangular grid without local cut vertices, but including an interior vertex, for which there is no through-edge Hamiltonian path.




Fig. 13. Example of a conforming quadrilateral grid without local cut vertices, but including an interior vertex, for which there is no Hamiltonian path.

Counterexample 6 *There exists a conforming quadrilateral grid with no local cut vertices and at least one interior vertex for which there is no Hamiltonian path. See Fig. 13.*

The proofs of Theorems 2 and 3 break down when applied to quadrilaterals and hexahedra because, in the notation of those theorems, E_{k+1} is not guaranteed to contain either the in-vertex or out-vertex of E_i . Without that condition, it is more difficult, and in some cases impossible, to modify the cycle in a way that adds E_{k+1} to it.

However, there are some other transformations of the cycle that can be applied to insert E_{k+1} into the cycle in some situations. These include some situations where

the grid is 1-nonconforming. Most of these transformations apply not only to quadrilaterals and hexahedra, but to any shape of element, including triangles and tetrahedra, and to mixed elements. Thus one could write a general program that applies to any grid. The illustrations will show the application of the transformations to quadrilaterals. They show examples of the transformations, but are not exhaustive.

An algorithm that attempts to find a through-vertex Hamiltonian cycle begins by picking two elements that share an edge (face in three dimensions) and constructing a cycle consisting of these two elements and two of the vertices they share. Then repeatedly pick an element that is not in the cycle but shares an edge (face in three dimensions) with an element in the cycle, and attempt to add it to the cycle by trying all of the following transformations that apply until one succeeds. If none of them succeed then try another element and come back to this one later. If there are elements not yet added to the cycle and none of the transformations work with any of the remaining elements, then you must insert a “broken link” creating a discontinuity in the path, i.e., insert an element such that either it is not adjacent to the previous or next element in the cycle, or the in-vertex and out-vertex do not match. When all elements have been added to the (possibly broken) cycle, the algorithm finishes.

In the following transformations, the cycle contains the segment $v_{i-2}E_{i-1}v_{i-1}E_i v_i E_{i+1}v_{i+1}$ and E is an element that shares an edge (face in three dimensions) with E_i and is not in the cycle.

Transformation 1 If E contains v_i and shares another vertex, $u \neq v_{i-1}$, with E_p , then replace the segment with $v_{i-2}E_{i-1}v_{i-1}E_i u E_p v_i E_{i+1}v_{i+1}$. See Fig. 14.

This is the same transformation that was used in the proof of Theorem 2.




Fig. 14. Example of Transformation 1.

Transformation 2 If E contains v_{i-1} and shares another vertex, $u \neq v_p$ with E_p , then replace the segment with $v_{i-2}E_{i-1}v_{i-1}Eu v_i E_{i+1}v_{i+1}$.

This is like the previous transformation, but using the in-vertex instead of the out-vertex.

Transformation 3 If E contains both v_{i-1} and v_p and E_{i-1} shares another vertex, $u \neq v_{i-2}$ with E_p , then replace the segment with $v_{i-2}E_{i-1}u E_i v_i E_{i+1}v_{i+1}$. See Fig. 15.




Fig. 15. Example of Transformation 3.

Transformation 4 If E contains both v_{i-1} and v_p and E_{i+1} shares another vertex, $u \neq v_{i+1}$ with E_p , then replace the segment with $v_{i-2}E_{i-1}v_{i-1}Ev_i u E_{i+1}v_{i+1}$.

This is like the previous transformation, but with the change occurring at the out-vertex of E_i instead of the in-vertex.

Transformation 5 If E shares a vertex $u_1 \neq v_i$ with E_p and shares a vertex u_2 with E_{i-1} , $u_2 \neq u_1$ and $u_2 \neq v_{i-2}$, then replace the segment with $v_{i-2}E_{i-1}u_2 E u_1 E_i v_i E_{i+1}v_{i+1}$. See Fig. 16.




Fig. 16. Example of Transformation 5.

This transformation can also handle some forms of hanging nodes, as shown in Fig. 17.




Fig. 17. Example of Transformation 5 with hanging node.

Transformation 6 If E shares a vertex $u_1 \neq v_{i-1}$ with E_p and shares a vertex u_2 with E_{i+1} , $u_2 \neq u_1$ and $u_2 \neq v_{i+1}$, then replace the segment with $v_{i-2}E_{i-1}v_{i-1}E_iu_1Eu_2E_{i+1}v_{i+1}$.

This is like the previous transformation, but with the changes occurring at E_{i+1} instead of E_{i-1} .

Transformation 7 If E contains neither v_{i-1} nor v_p but it shares a vertex u_1 with E_{i-1} , $u_1 \neq v_{i-2}$, and shares a vertex u_2 with E_p , $u_2 \neq u_1$, then replace the segment with $v_{i-2}E_{i-1}u_1Eu_2E_i v_i E_{i+1}v_{i+1}$. See Fig. 18.




Fig. 18. Example of Transformation 7.

Transformation 8 If E contains neither v_{i-1} nor v_p but it shares a vertex u_1 with E_{i+1} , $u_1 \neq v_{i+1}$, and shares a vertex u_2 with E_p , $u_2 \neq u_1$, then replace the segment with $v_{i-2}E_{i-1}v_{i-1}E_iu_2Eu_1E_{i+1}v_{i+1}$.

This is like the previous transformation, but with the changes occurring at E_{i+1} instead of E_{i-1} .

Transformation 9 If E contains v_i and there is another element, F , that is not on the cycle, shares a vertex, $u_1 \neq v_p$ with E , and shares a vertex u_2 with E_p , $u_2 \neq u_1$ and $u_2 \neq v_{i-1}$, then replace the segment with $v_{i-2}E_{i-1}v_{i-1}Eu_2Fu_1Ev_iE_{i+1}v_{i+1}$.

This transformation can handle some instances of hanging nodes, as shown in Fig. 19. It can also handle case 2 in the proof of Theorem 3, although it places the elements in a different order than that used in the proof.




Fig. 19. Example of Transformation 9.

Transformation 10 If E contains v_{i-1} and there is another element, F , that is not on the cycle, shares a vertex, $u_1 \neq v_{i-1}$, with E , and shares a vertex u_2 with E_p , $u_2 \neq u_1$ and $u_2 \neq v_p$, then replace the segment with $v_{i-2}E_{i-1}v_{i-1}Eu_1Fu_2E_i v_i E_{i+1}v_{i+1}$.

This is like the previous transformation, but with E sharing the in-vertex of E_i instead of the out-vertex.

Transformation 11 If E contains both v_{i-1} and v_p , then replace the segment with $v_{i-2}E_{i-1}v_{i-1}Ev_iE_{i+1}v_{i+1}$.

This is the swapping of one element for another that was described in case 3 of the proof of Theorem 3. One would next attempt to add E_i back into the cycle. If done recursively, this will handle case 3b of Theorem 3.

5. Conclusion

We considered the existence of various types of Hamiltonian paths and cycles in two-dimensional grids consisting of triangles or quadrilaterals, and three-dimensional grids consisting of tetrahedra or hexahedra. The types of Hamiltonian paths and cycles are distinguished by whether passage from one element to the next must be associated with an edge (through-edge Hamiltonian), must be associated with a vertex (through-vertex Hamiltonian), or can pass through either (unconstrained Hamiltonian). The existence results presented in this paper can be summarized as:

1. There exists an unconstrained Hamiltonian path in any conforming triangular grid with no local cut vertices (previously known result). This is easily extended to Hamiltonian cycle.
2. There exist triangular grids for which there is no through-edge Hamiltonian path (previously known result). This also holds for through-face Hamiltonian paths in tetrahedral grids.
3. There exists a through-vertex Hamiltonian cycle (and hence through-vertex Hamiltonian path) for any conforming triangular grid with no local cut vertices and at least one interior vertex.
4. There exists a through-vertex Hamiltonian cycle (and hence through-vertex Hamiltonian path, unconstrained Hamiltonian cycle and unconstrained Hamiltonian path) for any tetrahedral grid that contains no local cut vertices and no local cut edges.
5. There exist rectangular grids for which there is no Hamiltonian path or cycle of any type. This also holds for hexahedral grids.

Examples were given to show that the stated conditions are essential. Some open questions remain:

1. It is not known if the absence of local cut vertices is essential for the existence of a Hamiltonian path in a triangular grid. (The absence of cut vertices is essential, and the absence of local cut vertices is essential for a Hamiltonian cycle.)
2. It is not known if the presence of an interior vertex is essential for the existence of a Hamiltonian path in a triangular grid. (It is essential for a Hamiltonian cycle.)
3. The conditions under which any type of Hamiltonian path or cycle exists in a rectangular or hexahedral grid are not known.

The existence proofs for through-vertex Hamiltonian cycles in triangular and tetrahedral grids are constructive proofs. An efficient algorithm to find such a cycle can be derived from the construction in the proofs. Another algorithm, which can be applied to elements of any shape, was outlined. This algorithm is not guaranteed to find a through-vertex Hamiltonian cycle even if one exists, but it is likely to produce a (possibly broken) cycle that is close.

6. References

- [1] J. R. Pilkington and S. B. Baden, Dynamic partitioning of non-uniform structured workloads with spacefilling curves, IEEE Trans. on Par. and Dist. Sys. 7(3), 288-300 (1996).
- [2] J. Flaherty, R. Loy, M. Shephard, B. Szymanski, J. Teresco, and L. Ziantz, Adaptive local refinement with octree load-balancing for the parallel solution of three-dimensional conservation laws, J. Parallel Distrib. Comput. 47, 139-152 (1998).
- [3] W. F. Mitchell, The k -way refinement-tree partitioning method for adaptive grids, in progress.
- [4] K. Devine, B. Hendrickson, E. Boman, M. St. John, C. Vaughan, and W. F. Mitchell, Zoltan: A dynamic load-balancing library for parallel applications, User's Guide, Sandia Technical Report SAND99-1377, Sandia National Laboratories, Albuquerque, NM (2000).
- [5] G. Heber, R. Biswas, and G. R. Gao, Self-avoiding walks over two-dimensional adaptive unstructured grids, NAS Technical Report, NAS-98-007, NASA Ames Research Center, Moffett Field, CA (1998).
- [6] U. Dogrusoz and M. S. Krishnamoorthy, Hamiltonian cycle problem for triangle graphs, Computer Science Technical Report TR95-7, Rensselaer Polytechnic Institute, Troy, NY (1995).

About the author: William F. Mitchell is a computer scientist in the Mathematical and Computational Mathematics Division of the NIST Information Technology Laboratory. The National Institute of Standards and Technology is an agency of the Technology Administration, U.S. Department of Commerce.